

**This Page Is Inserted by IFW Operations
and is not a part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- **BLACK BORDERS**
- **TEXT CUT OFF AT TOP, BOTTOM OR SIDES**
- **FADED TEXT**
- **ILLEGIBLE TEXT**
- **SKEWED/SLANTED IMAGES**
- **COLORED PHOTOS**
- **BLACK OR VERY BLACK AND WHITE DARK PHOTOS**
- **GRAY SCALE DOCUMENTS**

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

Exhibit "E"

CENTER FOR DRUG EVALUATION AND RESEARCH

Application Number: NDA 20062/S027

APPROVAL LETTER



DEPARTMENT OF HEALTH & HUMAN SERVICES

Public Health Service

Food and Drug Administration
Rockville MD 20857

AUG 24 1999

DR

NDA 20-062/S-027

Marion Merrell Dow (Europe) AG
as General Partner of
Carderm Capital L.P.
c/o Westbroke Limited
Attention: Mr. Carlos A. Austin
Richmond House
12 Par-la Ville Road
P.O. Box HM 1022
Hamilton HM DX
Bermuda

Dear Mr. Austin:

Please refer to your supplemental new drug application dated January 7, 1999, received January 11, 1999, submitted under section 505(b) of the Federal Food, Drug, and Cosmetic Act for Cardizem CD (diltiazem hydrochloride) 180, 240, 300 and 360 mg Capsules.

We acknowledge receipt of your submissions dated May 11, June 18, and July 27, 1999. Your submission of June 18, 1999 constituted a complete response to our May 7, 1999 action letter.

This supplemental new drug application provides for a new dosage strength, 360 mg Capsules. The formulation of this new capsule strength is slightly modified from the other approved dosage strength capsules. Final printed labeling has been revised to incorporate information regarding this new dosage strength. In addition, the How Supplied statement was revised to read as follows:

Store at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) [see USP Controlled Room Temperature].

We have completed the review of this supplemental application, as amended, and have concluded that adequate information has been presented to demonstrate that the drug product is safe and effective for use as recommended in the submitted final printed labeling (package insert and immediate container and carton label submission dated June 18, 1999). Accordingly, the supplemental application is approved effective on the date of this letter.

We remind you that you must comply with the requirements for an approved NDA set forth under 21 CFR 314.80 and 314.81.

NDA 20-062/S-027

Page 2

If you have any questions, please contact:

David Roeder
Regulatory Health Project Manager
(301) 594-5313

Sincerely,

RSI 8/22/21

Raymond J. Lipicky, M.D.
Director
Division of Cardio-Renal Drug Products
Office of Drug Evaluation I
Center for Drug Evaluation and Research

CENTER FOR DRUG EVALUATION AND RESEARCH

APPLICATION NUMBER: NDA 20062/S027

APPROVABLE LETTER



DEPARTMENT OF HEALTH & HUMAN SERVICES

Food and Drug Administration
Rockville MD 20857

NDA 20-062/S-027

MAY -7 1999

Marion Merrell Dow (Europe) AG
as General Partner of
Carderm Capital L.P.
c/o Westbroke Limited
Attention: Carlos A. Austin
Richmond House
12 Par-la Ville Road
P.O. Box HM 1022
Hamilton HM DX
Bermuda

Dear Mr. Austin:

Please refer to your supplemental new drug application dated January 7, 1999, received January 11, 1999, submitted under section 505(b) of the Federal Food, Drug, and Cosmetic Act for Cardizem CD (diltiazem HCl) 120, 180, 240 and 300 mg Capsules.

We acknowledge receipt of your submission dated March 5, 1999.

This supplement provides for a new dosage strength, 360 mg Capsules. The formulation of this new capsule strength is slightly modified from the other approved dosage strength capsules.

We have completed the review of this application, as amended, and it is approvable. Before this application may be approved, however, it will be necessary for you to submit final printed labeling (FPL) revised as follows:

The Storage Statement should be revised in the package insert and the container labels to read as follows:

Store at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) [see USP
Controlled Room Temperature].

In addition, all previous revisions as reflected in the most recently approved labeling must be included. To facilitate review of your submission, please provide a highlighted or marked-up copy that shows the changes that are being made.

Please note that stability data at the 12-month time point for the 360 mg strength capsule at 25°C/60% RH and at 30°C/60% RH should be submitted in support of a 24-month expiration date.

Validation of the regulatory methods has not been completed. At the present time, it is the policy of the Center not to withhold approval because the methods are being validated. Nevertheless, we expect your continued cooperation to resolve any problems that may be identified.

Please submit 20 copies of the final printed labeling, ten of which are individually mounted on heavy weight paper or similar material.

If additional information relating to the safety or effectiveness of this drug becomes available, revision of the labeling may be required.

Within 10 days after the date of this letter, you are required to amend the supplemental application, notify us of your intent to file an amendment, or follow one of your other options under 21 CFR 314.110. In the absence of any such action FDA may proceed to withdraw the application. Any amendment should respond to all the deficiencies listed. We will not process a partial reply as a major amendment nor will the review clock be reactivated until all deficiencies have been addressed.

This product may be considered to be misbranded under the Federal Food, Drug, and Cosmetic Act if it is marketed with this change prior to approval of this supplemental application.

If you have any questions, please contact:

Mr. David Roeder
Regulatory Health Project Manager
(301) 594-5313

Sincerely yours,

151 5/7/69
Raymond J. Lipicky, M.D.
Director
Division of Cardio-Renal Drug Products
Office of Drug Evaluation I
Center for Drug Evaluation and Research

CENTER FOR DRUG EVALUATION AND RESEARCH

APPLICATION NUMBER: NDA 20062/S027

FINAL PRINTED LABELING

ure, propanolol appears to be different from its brother since it dilates. If combination therapy is needed or withdrawn in conjunction with propanolol, an adjustment in the propanolol dose may be warranted. (See WARNINGS.)

Diastolic. A study in healthy volunteers has shown a significant increase in peak diastolic plasma levels (28%) and area under the curve (57%) after a 1-week course of treatment at 1200 mg per day and a single dose of celecoxib 50 mg. Celecoxib produced similar, nonsignificant increases. The effect may be mediated by celecoxib's known inhibition of vascular prostaglandin-PGE₂, the enzyme system responsible for the first-phase metabolism of diazepam. Patients currently receiving diazepam therapy should be closely monitored for a change in pharmacological effect when initiating and discontinuing therapy with celecoxib. An adjustment in the diazepam dose may be warranted.

Cyclosporine. A pharmacokinetic interaction between diltiazem and cyclosporine has been observed in studies involving renal and cardiac transplant patients. In renal cardiac transplant recipients, a cycle of cyclosporine dose ranging from 15% to 40% was necessary to maintain Cyclosporine blood concentrations similar to those seen prior to the addition of diltiazem. If these agents are to be administered concurrently, cyclosporine concentrations should be lowered, especially when diltiazem is initiated, adjusted, or discontinued.

167 ~~XXXXXXXXXX~~

C. Reproduction studies

Abstract

44. *Journal of the American Medical Association*, 2000; 283: 2639-2644.

Adverse Reactions	Cardizem CD (n=687)	Placebo (n=301)
Headache	5.4%	1.6%
Dizziness	3.0%	3.6%
Nausea	3.3%	1.3%
Art Back		
Pain Chest	2.3%	0.5%
Edema	2.6%	1.3%
ECG		
Abnormality	1.6%	2.3%
Arthralgia	1.8%	1.7%

10-10-68

... ..

for a standard dose of 400

[illegible]

**Current Focus: Adolescent metabolic
syndrome (nonalcoholic fatty liver, etc.)**

been studied in clinical trials. The

10. Decays for the Treatment of

has an additive effect on the

--	--	--	--	--

[illegible]

001033

--	--	--	--	--

Strength		Quantity
250 mg	30 ml	
100 mg	30 ml	
50 mg	100 (100's)	
250 mg	30 ml	
100 mg	30 ml	
50 mg	100 (100's)	

Labeling: ORIGINAL
NDA No: 20-168 Rc'd. 6-21-99
Reviewed by: B. Smith 8-24-99

DATE: 6/18/99	CICERO JOB NO.: 13571
PRODUCT: LABEL, CARDIZEM CD, CAPSULES 360 MG, 90 CT.	
PRINTER: NEW JERSEY PACKAGING	
FIG 201	FIG 202





APPROVALS

FUNCTIONAL

PROOFREADING

REGULATORY

APPROVED
AUG 24 1999


NDC 0088-1799-42	Cardizem® CD diltiazem HCl	360mg		ONCE-A-DAY DOSAGE		90 Capsules Wendat Marion Roussel	<p>Each CARDIZEM® CD capsule contains 360 mg of diltiazem hydrochloride. U.S. ONLY Dosage and Administration: Read package insert for prescribing information. Warnings: Keep out of reach of children. Pharmacists: Dispense in light-resistant, tight container with child-resistant closure. Store at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) (see USP Controlled Room Temperature). Avoid excessive humidity.</p> <p>©1998, Hoechst Marion Roussel, Inc. Hoechst Marion Roussel, Inc. Kansas City, MO 64137 USA www.jrn.com</p>			3 0088-1799-42 1

CENTER FOR DRUG EVALUATION AND RESEARCH

APPLICATION NUMBER: NDA 20062/S027

CHEMISTRY REVIEW(S)

JUL 28 1999

CHEMIST'S REVIEW		1. ORGANIZATION HFD-110	2. NDA Number 20-062
3. Name and Address of Applicant (City & State) Carderm Capital L.P. c/o Westbroke Limited Raymond House 12 Par La Ville Road Hamilton, HM 12 Bermuda		4. Supplement(s) Number(s) Date(s) SCP-027 6/18/99	
5. Drug Name CARDIZEM CD	6. Nonproprietary Name Diltiazem hydrochloride		8. Amendments & Other (reports, etc) - Dates Orig - 1/7/99 BC-3/5/99
7. Supplement Provides For: Response to letter of May 7, 1999 for S-027.			
9. Pharmacological Category Ca antagonist (hypertension)	10. How Dispensed <input checked="" type="checkbox"/> Rx <input type="checkbox"/> OTC		11. Related IND(s)/NDA(s)/DMP(s)
12. Dosage Form(s) Capsules, CD (controlled diffusion, once-a-day)	13. Potency(ies) 120, 180, 240 and 300 mg/capsule		
14. Chemical Name and Structure 1,5-Benzothiazepin-4(5H)one, 3-(acetyloxy)-5-[2-(dimethylamino)ethyl]-2,3-dihydro-2-(4-(dimethylphenyl)-, monohydrochloride, (+)-cis-		15. Records/Reports Current <input type="checkbox"/> Yes <input type="checkbox"/> No Reviewed <input type="checkbox"/> Yes <input type="checkbox"/> No	
16. Comments: As requested in the letter, final proofs for the bottle labels and final printed labeling for the package insert showing the changes to the storage statement are included. Updated (12 month) stability report is included. The statistical analysis shows that the projected shelf life of the 360 mg capsules is greater than 24 months is also included. Labels - storage statement has been corrected. Satisfactory.			
17. Conclusions and Recommendations: For the final printed labeling, Hoechst Marion Roussel needs to use larger font. Unacceptable.			
18. REVIEWER			
Name Danute G. Cunningham	Signature 		Date Completed June 29, 1999
Distribution: <input checked="" type="checkbox"/> Original Jacket <input type="checkbox"/> Reviewer <input type="checkbox"/> Division File <input type="checkbox"/> CSO			

20062S27.AM1

1348-99

MAY 6 1999

CHEMIST'S REVIEW		1. ORGANIZATION HFD-110		2. NDA Number 20-062	
3. Name and Address of Applicant (City & State) Carderm Capital L.P. c/o Westbroke Limited Raymond House 12 Par La Ville Road Hamilton, HM 12 Bermuda				4. Supplement(s) Number(s) Date(s) SCF-027 1/7/99	
5. Drug Name CARDIZEM CD		6. Nonproprietary Name Diltiazem hydrochloride		8. Amendments & Other (reports, etc) - Dates B: 5/5/99 etc	
7. Supplement Provides For: strength addition of 360 mg capsule (slightly modified) to the Cardizem CD capsules.					
9. Pharmacological Category Ca antagonist (hypertension)		10. How Dispensed <input checked="" type="checkbox"/> Rx <input type="checkbox"/> OTC		11. Related IND(s)/NDA(s)/DMF(s) B.	
12. Dosage Form(s) Capsules, CD (controlled diffusion, once-a-day)		13. Potency(ies) 120, 180, 240 and 300 mg/capsule			
14. Chemical Name and Structure 1,5-Benzothiazepin-4(5H)one, 3-(acetyloxy)-5-[2-(dimethylamino)ethyl]-2,3-dihydro-2-(4-(dimethylphenyl)-, monohydrochloride, (+)-cis-				15. Records/Reports Current <input type="checkbox"/> Yes <input type="checkbox"/> No Reviewed <input type="checkbox"/> Yes <input type="checkbox"/> No	
16. Comments: The formulation change is found in the Active Bead. All other aspects, including the sustained release coating, of the manufacturing process of the capsule remain unchanged from that which is currently approved. This new capsule will be manufactured and controlled by Hoechst Marion Roussel, Inc. in Kansas City, MO. No change in formulation to the other approved strengths is proposed in this supplement.					
17. Conclusions and Recommendations: EES requested on 1/19/99. Acceptable on 5/5/99 Biopharmaceutics review requested on 1/19/99. Approvable - 4/23/99. Approvable - due to labeling issues. Storage statement has to be modified: Store at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) [see USP Controlled Room Temperature]. Container label - storage statement should be changed.					
18. REVIEWER					
Name Danute G. Cunningham		Signature <i>DS</i>		Date Completed May 5, 1999	
Distribution: <input checked="" type="checkbox"/> Original Jacket <input type="checkbox"/> Reviewer <input type="checkbox"/> Division File <input type="checkbox"/> CSO					

20062S27.SUP

5/5/99

CENTER FOR DRUG EVALUATION AND RESEARCH

APPLICATION NUMBER: NDA 20062/S027

**CLINICAL PHARMACOLOGY AND
BIOPHARMACEUTICS REVIEW(S)**

APR 23 1999

duj

Clinical Pharmacology/Biopharmaceutics Review

NDA 20-062

Serial #: SCF-027

Compound #: Cardizem CD 360mg Capsules

Hoechst Marion Roussel

Submission Date: January 7, 1999

Reviewer: Thomas A. Parmelee, Pharm.D.

Type of Submission: Supplement for New Formulation Study Report-Cardizem CD 360mg Capsules- A Bioequivalence Study and a Food Effect Study

BACKGROUND

NDA-062 has been approved for Cardizem CD (diltiazem HCl) Capsules in the strengths of 120mg, 180mg, 240mg, and 300mg. The maximum daily dose for diltiazem extended release capsules is established at 360mg. A new 360mg capsule formulation has been developed, and is the topic of this supplemental submission. The new 360mg capsule contains a formulation that is slightly modified from the currently approved lower strength capsules. The formulation change is found in the active bead of the drug product.

Two studies were submitted to the Office of Clinical Pharmacology and Biopharmaceutics for review. These studies were designed to show that the new Cardizem CD 360mg capsule formulation is bioequivalent to two Cardizem CD 180mg marketed capsules. One study is a bioequivalence study comparing single-dose and multiple-dose administration of the new 360mg capsules to the marketed 180mg strength capsules. The second study examines the effect of food on the single-dose pharmacokinetics of the new 360mg diltiazem capsule formulation. These studies are summarized in Appendix 1 and Appendix 2, respectively.

RESULTS

It appears that both lots of the new 360mg capsule formulation are bioequivalent to the marketed 180mg capsule formulation in the single-dose comparisons in terms of both AUC (0-inf) and C_{max} for both parent diltiazem and N-desmethyl metabolite.

In the steady-state comparisons, bioequivalency is met in terms of AUC, ss and C_{max}, ss between treatments, and only fails the 80-125% rule for Treatment B (lot # RH9738) in terms of C_{min}, ss for parent diltiazem.

The 90% confidence intervals between the high-fat breakfast treatment and the fasted treatment were within the 80-125% rule for both AUC (0-inf) and Cmax when looking at parent diltiazem and the N-desmethyl metabolite. Food does not appear to significantly affect the PK parameters of either lot of 360mg diltiazem capsules.

COMMENTS

- 1) Gender should not have been considered inclusion/exclusion criteria for these two clinical studies unless there was a specific reason for doing so. This point was mentioned to the sponsor via a teleconference before the start of the study.
- 2) The dissolution specifications are appropriate for the new strength capsule.
- 3) The draft prescription labeling submitted from the sponsor shows that the 360mg capsules contain black iron oxide, FD&C Blue #1, and starch. These ingredients were not listed in the composition of the capsules for review.

RECOMMENDATIONS

The new dosage strength for Cardizem CD 360mg capsules is approvable from the standpoint of the Office of Clinical Pharmacology and Biopharmaceutics. The comment above regarding the draft prescription labeling was conveyed to the review chemist. The dissolution methodology and specifications for the new strength are:

Apparatus: USP Type 2 (paddle)
Speed: 100 rpm
Media: 900mL degassed 0.1N HCl
Temperature: 37 C +/- 0.5 C

<u>Time (hrs.)</u>	<u>Specifications (%)</u>
6 hours	%
12 hours	%
18 hours	%
24 hours	NLT %
30 hours	NLT %

The draft prescription labeling (updated October 1998) and label included with the submission are attached to this review. The labeling for all diltiazem products is currently being updated and reviewed by this division (updated November 1998). The labeling for this new Cardizem CD 360mg capsule formulation should reflect the final printed labeling decided upon by the sponsor and the Agency for all Cardizem CD products.

IS/
Thomas A. Parmelee, Pharm.D.

4/23/99

APPENDIX 1

"BIOEQUIVALENCE OF 360MG DILTIAZEM HCL FORMULATIONS AND CARDIZEM CD AFTER SINGLE AND MULTIPLE DOSE ADMINISTRATIONS IN HEALTHY MALE SUBJECTS"

STUDY: Protocol # DZPR0207
Report K-98-0235-D

SPONSOR: Licensed to:
Hoechst Marion Roussel Inc.
P.O. Box 9627, H3-M2112
Kansas City, MO 64134-0627

Authorized by:
Carderm Capital L.P.
Raymond House
12 Par La Ville Road
Hamilton, HM 12 Bermuda

INVESTIGATOR AND STUDY SITE:

OBJECTIVES:

To determine whether 360mg Diltiazem HCL capsule formulations are bioequivalent to marketed 180mg Cardizem CD capsules.

FORMULATIONS:

- 1) Diltiazem 360mg capsules (lot# RH9736); Batch size
- 2) Diltiazem 360mg capsules (lot# RH9738); Batch size
- 3) Cardizem CD 180mg marketed capsules (lot# P31048)

The following table shows the composition of the new formulation of Cardizem CD 360mg Capsules:

STUDY DESIGN:

The study design was a randomized, open-label, single- and multiple-dose, 3-period, crossover study with a washout period of 12 days between treatments. The study population was 26 healthy, non-smoking males between the ages of 18 to 45 years. Subjects received each of the three treatment regimens in a randomized fashion:

Treatment A: One diltiazem 360mg capsule (RH9736) given as a SD on day 1, and then q.d. on days 3-9.

Treatment B: One diltiazem 360mg capsule (RH9738) given as a SD on day 1, and then q.d. on days 3-9.

Treatment C: Two Cardizem CD 180mg capsules (P31048) given as a SD on day 1, and then q.d. on days 3-9.

Subjects were continuously monitored for general health and any adverse reactions. Heart rate, blood pressure, and 12-lead ECG recordings, clinical chemistry, and hematological exams were done before the study and upon completion. Plasma samples were collected before the SD on day 1 and 1, 2, 3, 4, 5, 6, 8, 10, 12, 14, 16, 18, 21, 24, 36, and 48 hours following the dose. The subjects received seven days of multiple dosing during days 3-9. Trough plasma samples were obtained before the dose on days 8 and 9, and 1, 2, 3, 4, 5, 6, 8, 10, 12, 14, 16, 18, 21, and 24 hours following the day 9 dose.

ASSAY:

Table B: Dissolution Data for Cardizem CD 360mg Capsules

Time	Specifications (%)	RH9738 (%)	Percent Dissolved RH9736 (%)	P31048 180mg (%)
3 hours				
6 hours	%			
9 hours				
12 hours	%			
15 hours				
18 hours	%			
24 hours	NLT %			
30 hours	NLT %			

DATA ANALYSIS:

Pharmacokinetic analysis of diltiazem and MA metabolite concentrations in plasma was conducted by non-compartmental methods. The metabolite DAD concentrations were presented by descriptive statistics only (mean, standard deviation, CV%). The primary PK comparisons include C_{max} and AUC (0-inf) following single dose administration; and C_{max,ss}, C_{min,ss}, trough plasma concentrations (days 8, 9, and 10), and AUC_{ss} for multiple dose steady-state findings.

Comparisons between treatments were made for diltiazem and MA metabolite pharmacokinetic parameters and trough plasma levels. An analysis of variance (ANOVA) was performed for each parameter using PROC MIXED SAS with terms for sequence, subject within sequence, period, and treatment. Least square means, treatment differences, and 90% confidence intervals for treatment differences were determined. These log-transformed results were back-transformed by exponentiation to obtain adjusted means, treatment ratios, and 90% confidence intervals for these treatment ratios. Each lot of the diltiazem 360mg (Treatments A and B) was compared to the marketed reference Cardizem CD 180mg (Treatment C). Bioequivalence was to

be concluded if the limits of the 90% confidence interval on the ratio of treatment means falls entirely within the 80-125% range.

Trough plasma concentrations for each treatment were also compared using an ANOVA with terms for subject and day. From this ANOVA, least square means for each day, estimated differences between days, and 90% confidence intervals for the differences between days were calculated. These log-transformed results were back-transformed by exponentiation to obtain adjusted means, day ratios, and 90% confidence intervals for these ratios.

RESULTS:

Both lots of the 360mg capsules appear to be bioequivalent to the reference Cardizem CD 180mg capsules in the single-dose comparisons. Treatment A (lot # RH9736) appears to be bioequivalent to the reference capsules in the multiple-dose steady state comparison, however, treatment B (lot # 9738) is outside the 80-125% BE limits for C_{min}, ss. Please refer to the following tables and figures:



Figure 3. Mean diltiazem plasma concentrations following 360 mg single dose of once-daily capsules on day 1, protocol D2P10307.

A - lot RH9736 based (n=24); B - lot 9738 based (n=23); C - lot P1043 based (n=23).
(Cardizem CD)



Figure 4. Mean diltiazem plasma concentrations following 360 mg dose times 6 of once-daily capsules on day 6, -24 hours = trough sample on day 6, Protocol D2P10307.

A - lot RH9736 based (n=24); B - lot 9738 based (n=23); C - lot P1043 based (n=23).
(Cardizem CD)




Figure 5. Mean N-desmethyldiazepam plasma concentrations following 360 mg single dose on day 1 of once-daily capsules. Protocol DZP11207

A: 12 P10736 tested (n=24); B: 12 P10736 tested (n=24); C: 12 P10736 tested (n=24)
(Continued on p. 22)




Figure 6. Mean N-desmethyldiazepam plasma concentrations following 360 mg dose (time 0) on day 2 of once-daily capsules. 24 hours = trough sample on day 2. Protocol DZP11207

A: 12 P10736 tested (n=24); B: 12 P10736 tested (n=24); C: 12 P10736 tested (n=24)
(Continued on p. 22)

30 October FINAL

Table 9. Mean diltiazem (DTZ) pharmacokinetic parameters following 360 mg single dose on day 1, Protocol DZPR0207

	TRT	Number	Raw mean	Adjusted mean	CV%	Pair	Ratio (%)	90% CI ^a on ratio	P value
AUC(0-∞) (ng/mL·h)	A	24	3437.58	3254.72	30.91	A/C	100.83	(88.5, 114.8)	0.916
	B	23	3676.08	3436.67	36.23	B/A	105.59	(92.7, 120.3)	0.487
	C	23	3478.85	3228.07	39.54	B/C	106.46	(93.4, 121.3)	0.425
C _{max} (ng/mL)	A	24	170.69	160.18	38.54	A/C	102.47	(90.6, 115.8)	0.740
	B	23	169.69	158.35	35.20	B/A	98.86	(87.4, 111.8)	0.876
	C	23	166.58	156.32	36.49	B/C	101.30	(89.6, 114.5)	0.861
t _{1/2} (h)	A	24	6.98	6.86	16.61	A/C	95.33	(86.7, 104.8)	0.403
	B	23	7.48	7.10	45.65	B/A	103.51	(94.1, 113.9)	0.546
	C	23	7.30	7.20	19.48	B/C	98.68	(89.7, 108.6)	0.816
t _{max} (h)	A	24	13.08	12.17	35.13	A/C	106.27	(88.0, 128.3)	0.589
	B	23	13.22	12.45	30.52	B/A	102.34	(84.8, 123.6)	0.837
	C	23	12.39	11.45	34.30	B/C	108.77	(90.0, 131.5)	0.460

^a percent ratio and 90% confidence interval (CI) were calculated from ANOVA using log transformed data

TRT A = one 360 mg CD capsule (lot RH9736) given fasted

TRT B = one 360 mg CD capsule (lot RH9738) given fasted

TRT C = two 180 mg Cardizem CD Capsule (lot P31048) given fasted

Supporting Data:

Appendix B.3.3 Details of treatment comparisons, diltiazem single dose pharmacokinetic parameters, page 214 and Appendix C.2.2 Pharmacokinetic listings, page 639

Table 10. Mean diltiazem (DTZ) steady-state pharmacokinetic parameters following 360 mg dose on day 9, protocol DZPR0207

	TRT	Number	Raw mean	Adjusted mean	CV%	Pair	Ratio (%)	90% CI ^b on ratio	P value
AUC _{ss} (ng/mLxh)	A	24	3754.53	3551.98	28.57	A/C	100.69	(94.0, 107.9)	0.868
	B	23	3896.27	3558.25	36.19	B/A	100.18	(93.5, 107.3)	0.966
	C	23	3811.93	3527.75	33.13	B/C	100.86	(94.2, 108.1)	0.834
C _{max,ss} (ng/mL)	A	24	224.18	212.39	29.46	A/C	89.55	(83.5, 96.1)	0.011
	B	23	245.21	225.41	34.48	B/A	106.13	(98.9, 113.9)	0.163
	C	23	256.14	237.17	33.81	B/C	95.04	(88.6, 101.9)	0.230
C _{min,ss} (ng/mL)	A	24	97.29	87.97	38.53	A/C	104.02	(92.8, 116.6)	0.564
	B	23	109.15	97.94	41.76	B/A	111.33	(99.3, 124.8)	0.122
	C	23	94.05	84.57	41.10	B/C	115.80	(103.3, 129.8)	0.036
RATIO (C _{max,ss} / C _{min,ss})	A	24	2.55	2.41	49.16	A/C	85.77	(77.1, 95.4)	0.020
	B	23	2.36	2.31	25.23	B/A	95.78	(86.1, 106.6)	0.501
	C	23	2.89	2.81	29.11	B/C	82.14	(73.8, 91.4)	0.004
t _{max} (h)	A	24	9.75	9.38	43.75	A/C	149.54	(127.7, 175.1)	<0.001
	B	23	6.65	6.30	35.86	B/A	67.15	(57.3, 78.7)	<0.001
	C	23	6.65	6.27	36.15	B/C	100.42	(85.8, 117.5)	0.965
Trough Plasma Conc ^a (ng/mL)	A	24	116.96	110.78	31.37	A/C	108.12	(100.8, 116.0)	0.069
	B	23	119.28	110.00	36.39	B/A	99.29	(92.5, 106.5)	0.866
	C	23	110.28	102.46	34.20	B/C	107.36	(100.1, 115.2)	0.097

^a mean of trough plasma concentrations on days 8, 9, and 10
^b percent ratio and 90% confidence interval (CI) were calculated from ANOVA using log transformed data
 TRT A = one 360 mg CD capsule (lot RH9736) given on days 3 through 9
 TRT B = one 360 mg CD capsule (lot RH9738) given on days 3 through 9
 TRT C = two 180 mg Cardizem CD Capsule (lot P31048) given on days 3 through 9
 Supporting Data:
 Appendix B.3.7 Details of treatment comparisons, diltiazem steady state pharmacokinetic parameters, page 223
 Appendix C.2.2 Pharmacokinetic listings, page 639

Table 11. Mean N-desmethyldiltiazem (MA) pharmacokinetic parameters following 360 mg dose on day 1, Protocol DZPR0207									
	TRT	Number	Raw mean	Adjusted mean	CV%	Pair	Ratio (%)	90% CI ^a on ratio	P value
AUC(0-∞) (ng/mLxh)	A	24	1246.89	1176.76	32.17	A/C	99.12	(87.2, 112.6)	0.907
	B	23	1402.22	1272.85	56.64	B/A	108.17	(95.2, 122.9)	0.309
	C	23	1263.94	1187.27	36.93	B/C	107.21	(94.3, 121.9)	0.866
C _{max} (ng/mL)	A	24	43.05	40.43	35.32	A/C	97.49	(87.5, 108.6)	0.695
	B	23	43.37	41.12	28.05	B/A	101.72	(91.3, 113.4)	0.792
	C	23	43.86	41.47	29.52	B/C	99.17	(89.0, 110.5)	0.898
T _{1/2} (h)	A	24	11.16	10.96	18.53	A/C	99.14	(86.2, 114.0)	0.917
	B	23	13.79	12.09	86.90	B/A	110.31	(95.9, 126.9)	0.245
	C	23	11.22	11.06	23.65	B/C	109.36	(95.0, 125.9)	0.291
T _{max} (h)	A	24	16.63	16.12	22.49	A/C	100.83	(88.2, 115.2)	0.918
	B	23	16.52	15.38	47.86	B/A	95.42	(83.5, 109.1)	0.559
	C	23	16.09	15.99	19.20	B/C	96.21	(84.1, 110.0)	0.631
^a percent ratio and 90% confidence interval (CI) were calculated from ANOVA using log transformed data TRT A = one 360 mg CD capsule (lot RH9736) given fasted TRT B = one 360 mg CD capsule (lot RH9738) given fasted TRT C = two 180 mg Cardizem CD Capsule (lot P31048) given fasted Supporting Data: Appendix B.3.14 Details of treatment comparisons, MA single dose pharmacokinetic parameters, page 237 Appendix C.2.2 Pharmacokinetic listings, page 639									

Table 12. Mean N-desmethyldiltiazem (MA) steady-state pharmacokinetic parameters following 360 mg dose on day 8, Protocol DZPR0207

	TRT	Number	Raw mean	Adjusted mean	CV%	Pair	Ratio (%)	90% CI ^b on ratio	P value
AUC _{ss} (ng/mLxh)	A	24	1333.87	1254.56	31.56	A/C	98.30	(94.1, 102.7)	0.512
	B	23	1344.84	1254.89	33.11	B/A	100.03	(95.7, 104.5)	0.992
	C	23	1365.51	1276.27	33.37	B/C	98.33	(94.1, 102.7)	0.517
C _{max,ss} (ng/mL)	A	24	70.41	66.52	31.41	A/C	97.52	(93.0, 102.2)	0.376
	B	23	68.45	64.15	30.65	B/A	96.45	(92.0, 101.1)	0.205
	C	23	72.68	68.21	32.45	B/C	94.05	(89.7, 98.6)	0.034
C _{min,ss} (ng/mL)	A	24	41.31	37.48	39.71	A/C	102.20	(95.1, 109.8)	0.614
	B	23	43.80	40.62	35.07	B/A	108.36	(100.8, 116.5)	0.068
	C	23	40.07	36.68	38.79	B/C	110.73	(103.1, 119.0)	0.022
RATIO (C _{max,ss} / C _{min,ss})	A	24	1.82	1.78	33.29	A/C	95.41	(89.5, 101.8)	0.227
	B	23	1.59	1.58	12.60	B/A	89.04	(83.5, 95.0)	0.004
	C	23	1.88	1.86	20.76	B/C	84.95	(79.6, 90.6)	<0.001
t _{max} (h)	A	24	13.54	11.90	31.40	A/C	115.57	(92.6, 144.2)	0.278
	B	23	10.00	9.11	43.06	B/A	76.55	(61.3, 95.6)	0.049
	C	23	11.26	10.30	36.73	B/C	88.47	(70.8, 110.6)	0.361
Trough Plasma Conc ^a (ng/mL)	A	24	47.21	44.67	32.31	A/C	105.95	(101.4, 110.7)	0.032
	B	23	46.66	43.85	33.66	B/A	98.18	(94.0, 102.6)	0.486
	C	23	44.84	42.16	33.08	B/C	104.02	(99.6, 108.7)	0.137

^a mean of trough plasma concentrations on days 8, 9, and 10

^b percent ratio and 90% confidence interval (CI) were calculated from ANOVA using log transformed data

TRT A = one 360 mg CD capsule (lot RH9736) given on days 3 through 9

TRT B = one 360 mg CD capsule (lot RH9738) given on days 3 through 9

TRT C = two 180 mg Cardizem CD Capsule (lot P31048) given on days 3 through 9

Supporting Data:

Appendix B.3.18 Details of treatment comparisons, MA steady state pharmacokinetic parameters, page 246 and Appendix C.2.2 Pharmacokinetic listings, page 639

APPENDIX 2

"EFFECT OF FOOD ON THE SINGLE-DOSE PHARMACOKINETICS OF DILTIAZEM HCl 360MG FORMULATIONS IN HEALTHY MALE SUBJECTS"

STUDY: Protocol # DZPR0208
Report K-98-0236-D

SPONSOR: Licensed to:
Hoechst Marion Roussel Inc.
P.O. Box 9627, H3-M2112
Kansas City, MO 64134-0627

Authorized by:
Carderm Capital L.P.
Raymond House
12 Par La Ville Road
Hamilton, HM 12 Bermuda

INVESTIGATOR AND STUDY SITE:

OBJECTIVES:

To determine the effects of a high-fat breakfast on the rate and extent of absorption of a single oral dose of 360mg diltiazem HCl capsule formulation.

FORMULATIONS:

- 1) Diltiazem HCl 360mg capsules (lot# RH9736)
- 2) Diltiazem HCl 360mg capsules (lot# RH9738)

STUDY DESIGN:

The study design was a randomized, open-label, single-dose 4-period, crossover study with a washout period of 7 days between treatments. The study population was 22 healthy, non-smoking males between the ages of 18 to 45 years. Subjects received each of the four treatment regimens in a randomized fashion:

Treatment A: One diltiazem 360mg capsule (RH9736) dosed under fasting conditions.

Treatment B: One diltiazem 360mg capsule (RH9736) dosed with a high-fat breakfast.
Treatment C: One diltiazem 360mg capsule (RH9738) dosed under fasting conditions.
Treatment D: One diltiazem 360mg capsule (RH9738) dosed with a high-fat breakfast.

Subjects were continuously monitored for general health and adverse events. Heart rate, blood pressure, and 12-lead ECG recordings, clinical chemistry, and hematological exams were done before the study and upon completion. Heart rate, blood pressure (5 minutes supine), and lead II ECG measurements were taken 4 hours following each single dose. Plasma samples were collected before each dose on day 1 and 1, 2, 3, 4, 5, 6, 8, 10, 12, 14, 16, 18, 21, 24, 36, and 48 hours following the dose.

ASSAY:

DATA ANALYSIS:

Pharmacokinetic analysis of diltiazem and MA metabolite concentrations in plasma was conducted by non-compartmental methods. The metabolite DAD concentrations were presented by descriptive statistics only (mean, standard deviation, CV%). The primary PK comparisons include C_{max} and AUC (0-inf) for plasma concentrations.

Comparisons between treatments were made for diltiazem and MA metabolite pharmacokinetic parameters. An analysis of variance (ANOVA) was performed for each parameter using PROC MIXED SAS with terms for sequence, subject within sequence, period, and treatment. Least square means, treatment differences, and 90% confidence intervals for treatment differences were determined. These log-transformed results were back-transformed by exponentiation to obtain adjusted means, treatment ratios, and 90% confidence intervals for these treatment ratios. Treatment B was compared to Treatment A with Treatment A serving as the reference, and Treatment D was compared to Treatment C with Treatment C as the reference treatment. Equivalence was defined as the limits of the 90% confidence interval on the ratio of treatment means falling entirely within 80% to 125%.

RESULTS:

Twenty subjects completed all four treatments. The differences in AUC (0-inf) and Cmax between the high-fat and fasting treatments were small. The 90% confidence intervals for the differences between treatments were within the limits of 80% to 125% using the fasted treatments as the references. Please refer to the following tables and figures:

**APPEARS THIS WAY
ON ORIGINAL**




Figure 3. Mean diltiazem plasma concentrations following 360 mg single dose of once-daily capsules, Protocol DZPR0208

A= lot RH9736 fasted (n=20), B= lot RH9736 fed (n=20), C= lot RH9738 fasted (n=20), D= lot RH9738 fed (n=21).

Supporting Data:
Appendix C.2.2 Pharmacokinetic listings,

page 473




Figure 4. Mean N-desmethyldiltiazem plasma concentrations following 360 mg single dose of once-daily capsules, Protocol DZPR0208

A= lot RH9736 fasted (n=20), B= lot RH9736 fed (n=20), C= lot RH9738 fasted (n=20), D= lot RH9738 fed (n=21).

Supporting Data:
Appendix C.2.2 Pharmacokinetic listings,

page 473




Figure 5. Mean DAD plasma concentrations, Protocol DZPR0208

A= lot RH9736 fasted, B= lot RH9736 fed, C= lot RH9738 fasted, D= lot RH9738 fed.

Supporting Data:
Appendix C.2.2 Pharmacokinetic listings.

page 473

Table 9. Mean diltiazem (DTZ) pharmacokinetic parameters, 360 mg single dose, protocol DZPR0208

	TR T	Number	Raw mean	Adjusted mean	CV%	Pair	Ratio (%)	90% CI ^a on ratio	P value
AUC(0-∞) (ng/mL·h)	A	20	3384.33	3106.43	29.03	—	—	—	—
	B	20	3517.62	3240.02	31.23	B/A	104.30	(95.1, 114.4)	0.451
	C	20	3214.98	2961.00	27.04	—	—	—	—
	D	21	3633.28	3272.02	47.82	D/C	110.50	(100.7, 121.2)	0.077
C _{max} (ng/mL)	A	20	160.48	149.61	30.10	—	—	—	—
	B	20	179.60	166.93	34.52	B/A	111.58	(101.8, 122.3)	0.051
	C	20	153.51	144.68	24.63	—	—	—	—
	D	21	174.18	159.05	43.74	D/C	109.93	(100.3, 120.5)	0.089
t _{1/2} (h)	A	20	6.87	6.68	16.06	—	—	—	—
	B	20	6.65	6.49	13.47	B/A	97.16	(92.7, 101.8)	0.306
	C	20	6.77	6.60	13.55	—	—	—	—
	D	21	6.49	6.41	16.21	D/C	97.03	(92.6, 101.7)	0.283
t _{max} (h)	A	20	11.40	10.15	45.93	—	—	—	—
	B	20	10.10	9.33	40.37	B/A	91.93	(73.3, 115.3)	0.536
	C	20	13.00	11.85	38.18	—	—	—	—
	D	21	10.48	9.21	57.24	D/C	77.73	(62.2, 97.2)	0.065

^a percent ratio and 90% confidence interval (CI) were calculated from ANOVA using log transformed data

TRT A = one 360 mg CD capsule (lot RH9736) given fasted

TRT B = one 360 mg CD capsule (lot RH9736) given with high-fat breakfast

TRT C = one 360 mg CD capsule (lot RH9736) given fasted

TRT D = one 360 mg CD capsule (lot RH9736) given with high-fat breakfast

Supporting Data:

Appendix B.3.3 Details of treatment comparisons, page 201 and Appendix C.2.2 Pharmacokinetic listings, page 473

Table 10. Mean N-desmethyldiltiazem (MA) pharmacokinetic parameters, 360 mg single dose, protocol DZPR0208

	TRT	Number	Raw mean	Adjusted mean	CV%	Pair	Ratio (%)	90% CI ^a on ratio	P value
AUC (0-∞) (ng/mL·h)	A	20	1161.61	1083.07	27.09	-	-	-	-
	B	20	1196.27	1133.09	23.05	-	-	-	-
	C	20	1081.72	1011.27	24.71	B/A	104.62	(97.7, 112.0)	0.272
	D	21	1166.22	1116.67	29.43	D/C	110.42	(103.2, 118.2)	0.018
C _{max} (ng/mL)	A	20	41.29	39.18	27.14	-	-	-	-
	B	20	43.22	41.68	25.04	-	-	-	-
	C	20	38.12	36.43	21.19	B/A	106.38	(99.8, 113.4)	0.109
	D	21	40.04	38.85	24.78	D/C	106.64	(100.1, 113.6)	0.095
t _{1/2} (h)	A	20	10.22	9.89	17.45	-	-	-	-
	B	20	10.32	10.01	17.03	-	-	-	-
	C	20	9.97	9.64	17.05	B/A	101.20	(96.3, 106.4)	0.689
	D	21	10.40	10.15	22.52	D/C	105.24	(100.1, 110.6)	0.091
t _{max} (h)	A	20	16.85	16.29	18.99	-	-	-	-
	B	20	12.45	11.55	35.97	-	-	-	-
	C	20	16.10	15.77	18.68	B/A	70.88	(58.6, 85.7)	0.004
	D	21	12.14	10.63	50.92	D/C	68.68	(56.9, 82.9)	0.001

^a percent ratio and 90% confidence interval (CI) were calculated from ANOVA using log transformed data

TRT A = one 360 mg CD capsule (lot RH9736) given fasted

TRT B = one 360 mg CD capsule (lot RH9736) given with high-fat breakfast

TRT C = one 360 mg CD capsule (lot RH9736) given fasted

TRT D = one 360 mg CD capsule (lot RH9736) given with high-fat breakfast

Supporting Data:

Appendix B.3.7 Details of treatment comparisons, page 209 and Appendix C.2.2 Pharmacokinetic listings, page 473

CENTER FOR DRUG EVALUATION AND RESEARCH

APPLICATION NUMBER: NDA 20062/S027

ADMINISTRATIVE DOCUMENTS

RHPM Review of Final Printed Labeling

AUG 24 1999

Application: NDA 20-062
Cardizem CD (diltiazem HCl) Capsules

Applicant: Carderm Capital L.P.

Supplement Date: January 7, 1999

FPL Letter Date: June 18, 1999

FPL Receipt Date: June 21, 1999

Background

NDA 20-062/S-027 provides for a new dosage strength, 360 mg Capsules. The formulation of this new capsule strength is slightly modified from the other approved dosage strength capsules. An approvable letter was issued on May 7, 1999. In addition to the labeling changes under **DESCRIPTION** and **HOW SUPPLIED** relating to the new dosage strength, the approvable letter requested a revision of the **Storage Statement**.

Review

The applicant submitted final printed labeling in a submission dated June 18, 1999. The labeling was revised to include information on the 360 mg capsule under **DESCRIPTION** and **HOW SUPPLIED**. In addition, the **Storage Statement** was revised to read as follows:

Store at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) [see USP Controlled Room Temperature].

These changes were made in accordance with the requests in the approvable letter. An approval letter will be drafted for Dr. Lipicky's signature.


David Roeder
Regulatory Health Project Manager

cc: NDA 20-062
HFD-110
HFD-110/DRoeder/ABlount

CENTER FOR DRUG EVALUATION AND RESEARCH

APPLICATION NUMBER: NDA 20062/S027

CORRESPONDENCE

JUL 19 1999

Hoechst Marion Roussel, Inc.
Attention: Janet K. DeLeon
10236 Marion Park Drive
P.O. Box 9627
Kansas City, MO 64134-0627

Dear Ms. DeLeon:

Please refer to your January 7, 1999 supplemental new drug application (NDA) submitted under section 505(b) of the Federal Food, Drug and Cosmetic Act for CARDIZEM CD (diltiazem hydrochloride) Capsules, 180 mg, 240 mg, 300 mg and 360 mg.

The supplemental application provides for a new dosage strength, 360 mg Capsules. The formulation of this new capsule strength is slightly modified from the other approved dosage strength capsules.

We have completed our validation of the analytical methods for the 360 mg capsules and request the following additional information regarding the dissolution test:

The method refers to dissolution software used to correct for UV absorbance interference from diethyl phthalate, an excipient in the product. Attempts on April 8, 1999 by the analyst to get detailed information and explicit calculation formulas from your firm for dissolution calculations for the excipient contribution were not entirely successful. Please include a detailed description of the software and the calculations used to obtain the final results in the method.

The method does not specify whether aliquots taken out are replaced or not. If not replaced, please state whether final results are corrected for the volume taken during sampling. The validating analyst did not replace aliquots and corrected the volume withdrawn. It may be that sample aliquots are circulated back into the dissolution bath after samples are read. If this is the case, it should be stated in the method.

We would appreciate your prompt written response.

If you have any questions, please contact Danute G. Cunningham at (301) 594-5351 or Kasturi Srinivasachar, Ph.D. at (301) 594-5376.

Sincerely yours,

JSI 7-19-99
Kasturi Srinivasachar, Ph.D.
Chemistry Team Leader, DNDC I, for the
Division of Cardio-Renal Drug Products, (HFD-110)
Office of New Drug Chemistry
Center for Drug Evaluation and Research